

THE FUNDAMENTAL IDEAL AND π_2 OF HIGHER DIMENSIONAL KNOTS

S. J. LOMONACO, JR.

ABSTRACT. Let $(S^4, k(S^2))$ be a knot formed by spinning a polyhedral arc α about the standard 2-sphere S^2 in the 3-sphere S^3 . Then the second homotopy group of $S^4 - k(S^2)$ as a $Z\pi_1$ -module is isomorphic to each of the following:

- (1) The fundamental ideal modulo the left ideal generated by $a-1$, where a is the image in $\pi_1(S^4 - k(S^2))$ of a generator of $\pi_1(S^3 - \alpha)$.
- (2) The first homology group of the kernel of $\pi_1(S^3 - k(S^2)) \rightarrow \pi_1(S^4 - k(S^2))$

0. Introduction. A presentation of the second homotopy group of an arbitrary spun knot [3] was calculated as a $Z\pi_1$ -module in [4], [5], [6]. Professor John Milnor has conjectured that this is a presentation of the fundamental ideal modulo a suitably chosen element. In this paper, we show that this is actually the case. In particular,

THEOREM 2. *Let $k(S^2)$ be a 2-sphere formed by spinning an arc α about the standard 2-sphere S^2 in the 3-sphere S^3 . Let a denote the image in $\pi_1(S^4 - k(S^2))$ of a generator of $\pi_1(S^3 - \alpha)$ and let \mathfrak{I} be the fundamental ideal in $Z\pi_1$, i.e., the two sided ideal generated by all elements of the form $g-1$ ($g \in \pi_1$). Then the second homotopy group $\pi_2(S^4 - k(S^2))$ as a $Z\pi_1$ -module is $\mathfrak{I}/(a-1)$, where $(a-1)$ denotes the left ideal generated by $a-1$. The action of $Z\pi_1$ on π_2 is that induced by left multiplication on \mathfrak{I} .*

Moreover, we have yet another characterization of π_2 , namely

THEOREM 1. *Let $(S^4, k(S^2))$ be defined as in Theorem 2 above. Then $\pi_2(S^4 - k(S^2))$ as a $Z\pi_1$ -module is the first homology group of the kernel of $\pi_1(S^3 - k(S^2)) \rightarrow \pi_1(S^4 - k(S^2))$.*

I. Definition of a spun knot. Let S^2 be a standard 2-sphere in the 3-sphere S^3 and let α be a polyhedral arc with endpoints lying on S^2 and with interior lying entirely within one of the two components of $S^3 - S^2$.

Received by the editors June 5, 1972.

AMS (MOS) subject classifications (1970). Primary 57C45; Secondary 55E99, 55E15.

Key words and phrases. Spun knot, homotopy groups, homology groups, integral group ring, fundamental ideal, universal covering, fibration, Mayer-Vietoris sequence, Eilenberg-Mac Lane space.

© American Mathematical Society 1973

Generate a knotted 2-sphere $k(S^2)$ in S^4 by spinning α about S^2 while holding S^2 fixed. (For details see [3], [7].)

II. Proofs of theorems. Let $X = S^4 - k(S^2)$ and let $X_0 = S^3 - k(S^2)$ be the 3-dimensional cross-section of the knot. Let X_+ and X_- denote the closures of the two components of $X - X_0$. Similarly, let $X_{00} = S^2 - k(S^2)$ and X_{0+} and X_{0-} be the closures of the two components of $X_0 - X_{00}$. Let \tilde{X} be the universal covering of X , and $\tilde{X}_\pm, \tilde{X}_0, \tilde{X}_{0\pm}$, and \tilde{X}_{00} be the respective lifts of $X_\pm, X_0, X_{0\pm}, X_{00}$ to \tilde{X} .

The lifts $\tilde{X}_\pm, \tilde{X}_0, \tilde{X}_{0\pm}$ are all connected. This can be seen by inspecting the homotopy sequence for the fibration

$$\pi_1(X) \rightarrow \tilde{X}_i \rightarrow X_i$$

and noting that $\pi_1(X_i) \rightarrow \pi_1(X)$ is onto for $i = +, -, 0, 0+, 0-$. Moreover, since $\pi_1(X_i) \rightarrow \pi_1(X)$ for $i = +, -, 0+, 0-$ is an isomorphism onto [3], $\tilde{X}_\pm, \tilde{X}_{0\pm}$ are all simply connected.

Since \tilde{X}_\pm collapses to $\tilde{X}_{0\pm}$ via a deformation arising from the spinning, Hurewicz's theorem coupled with the asphericity of knots [2] gives $H_n(\tilde{X}_\pm) = 0$ for $n \geq 1$. Hence, from the Mayer-Vietoris sequence for the triad $(\tilde{X}; \tilde{X}_+, \tilde{X}_-)$, we have

$$H_2(\tilde{X}) \cong H_1(\tilde{X}_0).$$

But by the asphericity of knots [2], \tilde{X}_0 is the Eilenberg-Mac Lane space $K(\pi_1(\tilde{X}_0), 1)$. Inspecting the homology sequence for the fibration $\pi_1(\tilde{X}) \rightarrow \tilde{X}_0 \rightarrow X_0$, we have

THEOREM 1. *The second homotopy group $\pi_2(S^4 - k(S^2))$ as a $Z\pi_1$ -module is the first homology group of the kernel of $\pi_1(S^3 - k(S^2)) \rightarrow \pi_1(S^4 - k(S^2))$.*

Theorem 2 now follows from a close inspection of the exact sequence

$$0 \rightarrow H_1(\tilde{X}_0) \rightarrow H_0(\tilde{X}_{00}) \rightarrow H_0(\tilde{X}_{0+}) \oplus H_0(\tilde{X}_{0-})$$

arising from the Mayer-Vietoris sequence for $(\tilde{X}_0; \tilde{X}_{0+}, \tilde{X}_{0-})$.

Note that there is a deformation retraction of X_{0+} onto a 2-dimensional CW-complex K_{0+} which induces on X_{00} a deformation retraction onto a 1-dimensional CW-complex K_{00} . The CW-complex K_{00} consists of a single 1-simplex ξ_0 and a single vertex p . The element a of $\pi_1(X)$ carried by ξ_0 is the image of a generator of $\pi_1(X_{00})$. (For details see the lemma of [6].) Thus, \tilde{X}_{00} collapses to

$$\tilde{K}_{00} = \bigcup_{g \in \pi_1} g \xi_0.$$

By direct computation, $Z_0(\tilde{K}_{00}) = Z\pi_1 p$ and $B_0(\tilde{K}_{00}) = (a-1)p$, where

$(a-1)$ is the left ideal generated by $a-1$. Hence, $H_0(\tilde{X}_{00}) \cong Z\pi_1/(a-1)$ and

$$0 \rightarrow H_1(\tilde{X}_0) \rightarrow Z\pi_1/(a-1) \xrightarrow{\sigma_*} J \oplus J.$$

Again by direct computation, $\sigma_* = \varepsilon \oplus (-\varepsilon)$, where $\varepsilon: Z\pi_1/(a-1) \rightarrow J$ is the projection of the trivializer [1] (also called the augmentation [8]). Thus, $\text{Ker}(\sigma_*) = \text{Ker}(\varepsilon) \cap \text{Ker}(-\varepsilon) = \mathfrak{I}/(a-1)$ and

$$\mathfrak{I}/(a-1) \cong H_1(\tilde{X}_0) \cong H_2(\tilde{X}) \cong \pi_2(X).$$

REFERENCES

1. R. H. Crowell and R. H. Fox, *Introduction to knot theory*, Ginn, Boston, Mass., 1963. MR 26 #4348.
2. Ch. D. Papakyriakopoulos, *On Dehn's lemma and the asphericity of knots*, Ann. of Math. (2) 66 (1957), 1-26. MR 19, 761.
3. Emil Artin, *Zur isotopie zweidimensionalen Flächen im R^4* , Abh. Math. Sem. Univ. Hamburg, 4 (1925), 174-177.
4. J. J. Andrews and S. J. Lomonaco, *The second homotopy group of 2-spheres in 4-space*, Bull. Amer. Math. Soc. 75 (1969), 169-171. MR 38 #3863.
5. J. J. Andrews and S. J. Lomonaco, *The second homotopy group of 2-spheres in 4-space*, Ann. of Math. (2) 90 (1969), 199-204. MR 40 #3547.
6. S. J. Lomonaco, Jr., *The second homotopy group of a spun knot*, Topology 8 (1969), 95-98. MR 38 #6594.
7. ———, *The third homotopy group of higher dimensional knots* (to appear).
8. S. Mac Lane, *Homology*, Die Grundlehren der math. Wissenschaften, Band 114, Academic Press, New York; Springer-Verlag, Berlin, 1963. MR 28 #122.
9. Sze-Tsen Hu, *Homotopy theory*, Pure and Appl. Math., vol. 8, Academic Press, New York, 1959. MR 21 #5186.

STATE UNIVERSITY OF NEW YORK AT ALBANY, ALBANY, NEW YORK 12203

Current address (S. J. Lomonaco, Jr.): 44 Blue Spruce Lane, Ballston Lake, New York 12019